



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Linear Algebra and its Applications 426 (2007) 667–673

LINEAR ALGEBRA
AND ITS
APPLICATIONSwww.elsevier.com/locate/laa

An upper bound for $\|A^{-1}\|_{\infty}$ of strictly diagonally dominant M -matrices

Guang-Hui Cheng^{*}, Ting-Zhu Huang¹*School of Applied Mathematics, University of Electronic Science and Technology of China,
Chengdu, Sichuan 610054, PR China*

Received 9 February 2007; accepted 5 June 2007

Available online 16 June 2007

Submitted by M. Tsatsomeros

Abstract

Let A be a real strictly diagonally dominant M -matrix. We give a sharp upper bound for $\|A^{-1}\|_{\infty}$. Furthermore, the lower bound of the smallest eigenvalue $q(A)$ is established.

© 2007 Elsevier Inc. All rights reserved.

AMS classification: 15A42; 15A45; 65F15

Keywords: Diagonal dominance; M -matrix; Inverse M -matrix

1. Introduction

A square $n \times n$ matrix A is called a nonsingular M -matrix if there exists an $n \times n$ nonnegative matrix P such that

$$A = sI - P, \quad (1)$$

where I is the identity matrix, and $s > \rho(P)$, the spectral radius of nonnegative matrix P . If A is an M -matrix, there exists a positive eigenvalue of A equal to $\frac{1}{\rho(A^{-1})}$, where $\rho(A^{-1})$ is the Perron eigenvalue of the nonnegative matrix A^{-1} . We denote this eigenvalue by $q(A)$, then $q(A) = s - \rho(P)$ is also the minimum of the real parts of the eigenvalues of A . See [1–3], for example, for further discussion of this issue.

^{*} Corresponding author.

E-mail addresses: ghcheng@uestc.edu.cn (G.-H. Cheng), tzh Huang@uestc.edu.cn (T.-Z. Huang).

¹ Supported by NCET of China (NCET-04-0893).

Shivakumar et al. [4] gave the following results on two-sided bounds related to weakly chained diagonally dominant M -matrices.

Theorem 1.1 [4]. *Let $A = (a_{ij})$ be an $n \times n$ weakly chained diagonally dominant M -matrix, let $A^{-1} = (\alpha_{ij})$, and let $q = q(A)$, $N = \{1, 2, \dots, n\}$. Then*

$$q \leq \min\{a_{ii} : i \in N\}, \quad (2)$$

$$q \leq \max \left\{ \sum_{j \in N} a_{ij} : i \in N \right\}, \quad (3)$$

$$q \geq \min \left\{ \sum_{j \in N} a_{ij} : i \in N \right\}, \quad (4)$$

$$\frac{1}{M} \leq q \leq \frac{1}{m}, \quad (5)$$

where

$$M = \max_{i \in N} \sum_{j \in N} \alpha_{ij} = \|A^{-1}\|_{\infty} \quad \text{and} \quad m = \min_{i \in N} \sum_{j \in N} \alpha_{ij}.$$

Varah [7] gave the following result.

Theorem 1.2 [7]. *If $A = a_{ij} \in \mathbb{R}^{n \times n}$ is a strictly diagonally dominant matrix, then*

$$\|A^{-1}\|_{\infty} \leq \frac{1}{\min_i \left\{ |a_{ii}| - \sum_{j \neq i} |a_{ij}| \right\}}, \quad i \in N. \quad (6)$$

Remark 1.1. If the diagonal dominance of A is weak, i.e., $\min_i \{|a_{ii}| - \sum_{j \neq i} |a_{ij}|\}$ is small, then using Theorem 1.2 in estimating $\|A^{-1}\|_{\infty}$, the bound may yield a large value.

In this paper, the upper bound of $\|A^{-1}\|_{\infty}$ is improved, furthermore, by using (5), a new lower bound of $q(A)$ is obtained.

The paper is organized as follows. In Section 2, we present notations and some preliminary results. The main results are driven for M -matrices in Section 3.

2. Notations and preliminaries

In the following, we need the following definitions and results. They will be useful in the following proofs. For convenience, for any positive integer n , N denotes the set $\{1, 2, \dots, n\}$ throughout. Let $A = (a_{ij})$ be an $n \times n$ matrix. For any $i \in N$, denote

$$d_i = \frac{1}{|a_{ii}|} \sum_{j \neq i} |a_{ij}|, \quad J(A) = \{i \in N : d_i < 1\},$$

$$u_i = \frac{1}{|a_{ii}|} \sum_{j=i+1}^n |a_{ij}|,$$

$$l_k = \max \left\{ \frac{\sum_{j \neq i+k-1, k \leq j \leq n} |a_{i+k-1, j}|}{|a_{i+k-1, i+k-1}|} : i = 1, 2, \dots, n - k + 1 \right\}.$$

Definition 2.1 [4]. $A \in \mathbb{R}^{n \times n}$ is weakly chained diagonally dominant if for all $i \in N$, $d_i \leq 1$ and $J(A) \neq \emptyset$, and for all $i \in N$, $i \notin J(A)$, there exist indices i_1, i_2, \dots, i_k in N with $a_{i_r, i_{r+1}} \neq 0$, $0 \leq r \leq k-1$, where $i_0 = i$ and $i_k \in J(A)$.

Definition 2.2 [8]. $A \in \mathbb{R}^{n \times n}$ is an L -matrix if for all $i, j \in N$ with $i \neq j$, $a_{ij} \leq 0$ and $a_{ii} > 0$.

Definition 2.3 [8]. Let $A \in \mathbb{R}^{n \times n}$. A is strictly diagonally dominant if $J(A) = N$.

Lemma 2.1 [5]. A weakly chained diagonally dominant L -matrix is a nonsingular M -matrix.

We will denote by $A^{(n_1, n_2)}$ as the principal submatrix of A formed from all rows and all columns with indices between n_1 and n_2 inclusively; e.g., $A^{(2, n)}$ is the submatrix of A obtained by deleting the first row and the first column of A .

Lemma 2.2 [5]. Let A be an $n \times n$ weakly chained diagonally dominant M -matrix. Then $B = A^{(2, n)}$ is an $(n-1) \times (n-1)$ weakly chained diagonally dominant M -matrix, i.e., $B^{-1} = (\beta_{ij})$ exists and $\beta_{ij} \geq 0$, $i, j = 2, 3, \dots, n$.

Lemma 2.3 [5]. Let $A = (a_{ij})$ be a weakly chained diagonally dominant M -matrix and $A^{-1} = (\alpha_{ij})$. Then, for $i \neq j$,

$$\alpha_{ij} \leq d_i \alpha_{jj} \leq \alpha_{jj}. \quad (7)$$

Lemma 2.4 [6]. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be an M -matrix, and $A^{-1} = (\alpha_{ij})$. Then

$$\alpha_{ii} \geq \left(a_{ii} - \sum_{k \neq i} \frac{a_{ik} a_{ki}}{a_{kk}} \right)^{-1} \geq a_{ii}^{-1}, \quad i \in N. \quad (8)$$

3. Upper bounds for $\|A^{-1}\|_\infty$

In this section, we give upper bounds for the inverse of a strictly diagonally dominant M -matrix.

Theorem 3.1. Let $A = (a_{ij})$ be a weakly chained diagonally dominant M -matrix, $B = A^{(2, n)}$, $A^{-1} = (\alpha_{ij})_{i, j=1}^n$, and $B^{-1} = (\beta_{ij})_{i, j=2}^n$. Then, for $i, j = 2, \dots, n$,

$$\alpha_{11} = \frac{1}{\Delta}, \quad (9)$$

$$\alpha_{i1} = \frac{1}{\Delta} \sum_{k=2}^n \beta_{ik} (-a_{k1}), \quad (10)$$

$$\alpha_{1j} = \frac{1}{\Delta} \sum_{k=2}^n \beta_{kj}(-a_{1k}), \quad (11)$$

$$\alpha_{ij} = \beta_{ij} + \alpha_{1j} \sum_{k=2}^n \beta_{ik}(-a_{k1}), \quad (12)$$

where

$$\Delta = a_{11} - \sum_{k=2}^n a_{1k} \left(\sum_{i=2}^n \beta_{ki} a_{i1} \right) > 0. \quad (13)$$

Furthermore, if $J(A) = N$, we have

$$\Delta \geq a_{11}(1 - d_1 l_1) \geq a_{11}(1 - d_1). \quad (14)$$

Proof. Eqs. (9)–(13) were proved in [4]. For $k, j \geq 2$,

$$\sum_{i=2}^n \beta_{ki} a_{ij} = (B^{-1}B)_{kj} = \begin{cases} 1, & k = j, \\ 0, & k \neq j. \end{cases}$$

From (13), using Lemmas 2.3 and 2.4 we have

$$\begin{aligned} \Delta &= a_{11} - \sum_{k=2}^n a_{1k} \left(\sum_{i=2}^n \beta_{ki} a_{i1} \right) \\ &= a_{11} - \sum_{k=2}^n a_{1k} \left[\sum_{i=2}^n \beta_{ki} a_{ii} (1 - d_i) - \sum_{j=2}^n \sum_{i=2}^n \beta_{ki} a_{ij} \right] \\ &= a_{11} - \sum_{k=2}^n a_{1k} \left[\sum_{i=2}^n \beta_{ki} a_{ii} (1 - d_i) - 1 \right] \\ &= \sum_{k=1}^n a_{1k} + \sum_{k=2}^n (-a_{1k}) \sum_{i=2}^n \beta_{ki} a_{ii} (1 - d_i) \\ &\geq \sum_{k=1}^n a_{1k} + (1 - l_1) \sum_{k=2}^n (-a_{1k}) \sum_{i=2}^n \beta_{ki} a_{ii} \\ &\geq \sum_{k=1}^n a_{1k} + (1 - l_1) \sum_{k=2}^n (-a_{1k}) \sum_{i=2}^n |\beta_{ki} a_{ik}| \\ &= \sum_{k=1}^n a_{1k} + (1 - l_1) \sum_{k=2}^n (-a_{1k}) (2\beta_{kk} a_{kk} - 1) \\ &\geq \sum_{k=1}^n a_{1k} + (1 - l_1) \sum_{k=2}^n (-a_{1k}) \\ &= a_{11}(1 - d_1 l_1) \\ &\geq a_{11}(1 - d_1) \\ &\geq 0. \quad \square \end{aligned} \quad (15)$$

Corollary 3.2. If $A = (a_{ij})$ is an $n \times n$ row strictly diagonally dominant M -matrix, then $\Delta \geq a_{11}(1 - d_1 l_1) > a_{11}(1 - d_1) > 0$.

Proof. Since $A = (a_{ij})$ is an $n \times n$ strictly diagonally dominant M -matrix, $0 < d_i < 1, i \in N$ and $0 \leq l_1 < 1$, i.e., $a_{11}(1 - d_1 l_1) > a_{11}(1 - d_1) > 0$, the conclusion follows. \square

Theorem 3.3. Let $A = (a_{ij})$ be an $n \times n$ strictly diagonally dominant M -matrix and $B = A^{(2,n)}$, $A^{-1} = (\alpha_{ij})_{i,j=1}^n$, and $B^{-1} = (\beta_{ij})_{i,j=2}^n$. Then

$$\|A^{-1}\|_\infty \leq (1 - d_1)\alpha_{11} + \frac{d_1}{a_{11}(1 - d_1 l_1)} + \frac{1}{1 - d_1 l_1} \|B^{-1}\|_\infty, \quad (16)$$

and

$$\|A^{-1}\|_\infty \leq \frac{1}{a_{11}(1 - d_1 l_1)} + \left(1 + \frac{d_1}{1 - d_1 l_1}\right) \|B^{-1}\|_\infty. \quad (17)$$

Proof. Let

$$r_i = \sum_{j=1}^n \alpha_{ij}, \quad M_A = \|A^{-1}\|_\infty, \quad M_B = \|B^{-1}\|_\infty.$$

Then

$$M_A = \max\{r_i : i \in N\} \quad \text{and} \quad M_B = \max \left\{ \sum_{j=2}^n \beta_{ij} : 2 \leq i \leq n \right\}.$$

By Theorem 3.1 and Corollary 3.2,

$$\begin{aligned} r_1 &= \alpha_{11} + \sum_{j=2}^n \alpha_{1j} \\ &= \frac{1}{\Delta} + \frac{1}{\Delta} \sum_{k=2}^n (-a_{1k}) \sum_{j=2}^n \beta_{kj} \\ &\leq \frac{1}{\Delta} + \frac{1}{\Delta} \sum_{k=2}^n (-a_{1k}) M_B \\ &\leq \frac{1 + a_{11} d_1 M_B}{a_{11}(1 - d_1 l_1)}. \end{aligned} \quad (18)$$

Let $2 \leq i \leq n$. Then, using (7) in (9) and (10),

$$\sum_{k=2}^n \beta_{ik} (-a_{k1}) \leq d_i < 1.$$

From (12), with $2 \leq j \leq n$, we have

$$\alpha_{ij} \leq \beta_{ij} + \alpha_{1j} d_i < \beta_{ij} + \alpha_{1j}. \quad (19)$$

Thus, for $2 \leq i \leq n$, we obtain

$$r_i = \alpha_{i1} + \sum_{j=2}^n \alpha_{ij}$$

$$\begin{aligned}
&\leq \alpha_{11} + \sum_{j=2}^n (\beta_{ij} + \alpha_{1j} d_i) \\
&\leq (1 - l_1) \alpha_{11} + l_1 r_1 + M_B,
\end{aligned} \tag{20}$$

or, we can obtain

$$r_i \leq r_1 + M_B. \tag{21}$$

According to (20) and (21), we have

$$M_A \leq (1 - l_1) \alpha_{11} + l_1 r_1 + M_B \quad \text{and} \quad M_A \leq r_1 + M_B,$$

respectively. Using (18), the results follow. \square

Theorem 3.4. Let $A = (a_{ij})$ be an $n \times n$ strictly diagonally dominant M -matrix. Then

$$\|A^{-1}\|_{\infty} \leq \frac{1}{a_{11}(1 - u_1 l_1)} + \sum_{i=2}^n \left[\frac{1}{a_{ii}(1 - u_i l_i)} \prod_{j=1}^{i-1} \left(1 + \frac{u_j}{1 - u_j l_j} \right) \right]. \tag{22}$$

Proof. Apply induction with respect to k to $A^{(k,n)}$, using (17). \square

Remark 3.1. Using (16), (17) and (22) in (5), we can obtain new bounds of the smallest eigenvalue $q(A)$.

Theorem 3.5. Let $A = (a_{ij})$ be an $n \times n$ strictly diagonally dominant M -matrix. Then the bound in (22) is sharper than that in Theorem 3.3 in [4], i.e.,

$$\frac{1}{a_{11}(1 - u_1 l_1)} + \sum_{i=2}^n \left[\frac{1}{a_{ii}(1 - u_i l_i)} \prod_{j=1}^{i-1} \left(1 + \frac{u_j}{1 - u_j l_j} \right) \right] < \sum_{i=1}^n \left[a_{ii} \prod_{j=1}^i (1 - u_i) \right]^{-1}. \tag{23}$$

Proof. Since A is a strictly diagonally dominant matrix, $0 \leq l_k < 1$ for all k . Consequently, we have $\frac{1}{a_{11}(1 - d_1 l_1)} < \frac{1}{a_{11}(1 - d_1)}$ and $(1 + \frac{d_1}{1 - d_1 l_1}) \|B^{-1}\|_{\infty} < \frac{1}{1 - d_1} \|B^{-1}\|_{\infty}$, i.e.,

$$\frac{1}{a_{11}(1 - d_1 l_1)} + \left(1 + \frac{d_1}{1 - d_1 l_1} \right) \|B^{-1}\|_{\infty} < \frac{1}{a_{11}(1 - d_1)} + \frac{1}{1 - d_1} \|B^{-1}\|_{\infty}. \tag{24}$$

The inequality (24) shows that the bound in (17) is better than that in Lemma 3.2 in [4], further, the bound in (22) is sharper than that in Theorem 3.3 in [4]. \square

In the following, we give a numerical example to illustrate the results obtained in Section 3.

Example 3.1. Let

$$A = \begin{bmatrix} 1 & 0 & -0.2 \\ -0.8 & 1 & -0.1 \\ -0.9 & 0 & 1 \end{bmatrix}.$$

By direct calculations with Matlab 7.0, we have

$$A^{-1} = \begin{bmatrix} 1.2195 & 0 & 0.2439 \\ 1.0854 & 1.0000 & 0.3171 \\ 1.0976 & 0 & 1.2195 \end{bmatrix}, \quad \|A^{-1}\|_{\infty} = 2.4025.$$

We have

$$\|A^{-1}\|_{\infty} \leq 10 \text{ (by Theorem 1.2),}$$

$$\|A^{-1}\|_{\infty} \leq 4.0278 \text{ (by Theorem 3.3 in [4]),}$$

$$\|A^{-1}\|_{\infty} \leq 3.8455 \text{ (by Theorem 3.4),}$$

respectively. It is obviously that the bound of Theorem 3.4 is sharper. Furthermore, we can use Theorem 3.4 and (5) in estimating $q(A)$.

Acknowledgments

The authors thank the anonymous referee for a thorough reading of the manuscript, useful comments, and pointing out several typos to improve the paper.

References

- [1] R. Horn, C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, New York, 1991.
- [2] C.R. Johnson, A Hadamard product involving M -matrices, *Linear and Multilinear Algebra* 4 (1977) 261–264.
- [3] M. Fiedler, C.R. Johnson, T.L. Markham, M. Neumann, A trace inequality for M -matrices and the symmetrizability of a real matrix by a positive diagonal matrix, *Linear Algebra Appl.* 101 (1988) 1–8.
- [4] P.N. Shivakumar, Joseph J. Williams, Qiang Ye, Corneliu A. Marinov, On two-sided bounds related to weakly diagonally dominant M -matrices with application to digital circuit dynamics, *SIAM J. Matrix Anal. Appl.* 17 (2) (1996) 298–312.
- [5] P.N. Shivakumar, K.H. Chew, A sufficient condition for nonvanishing of determinants, *Proc. Amer. Math. Soc.* 43 (1974) 63–66.
- [6] H.B. Li, T.Z. Huang, S.Q. Shen, H. Li, Lower bounds for the minimum eigenvalue of Hadamard product of an M -matrix and its inverse, *Linear Algebra Appl.* 420 (2007) 235–247.
- [7] J.M. Varah, A lower bound for the smallest singular value of a matrix, *Linear Algebra Appl.* 11 (1975) 3–5.
- [8] A. Berman, R.J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, third ed., Academic Press, New York, 1994 (Reprinted by SIAM, Philadelphia).